

This document was prepared in conjunction with work accomplished under Contract No. DE-AC09-96SR18500 with the U.S. Department of Energy.

This work was prepared under an agreement with and funded by the U.S. Government. Neither the U. S. Government or its employees, nor any of its contractors, subcontractors or their employees, makes any express or implied: 1. warranty or assumes any legal liability for the accuracy, completeness, or for the use or results of such use of any information, product, or process disclosed; or 2. representation that such use or results of such use would not infringe privately owned rights; or 3. endorsement or recommendation of any specifically identified commercial product, process, or service. Any views and opinions of authors expressed in this work do not necessarily state or reflect those of the United States Government, or its contractors, or subcontractors.

The Dependence of the Diffusion Parameters on the Dissolved H Concentration in the Dilute Phase of Pd–H

Ted B. Flanagan, Da Wang,
Chemistry Department, University of Vermont, Burlington VT 05405,
Kirk Shanahan
Savannah River National Laboratory, Hydrogen Bldg. 999-2W, Aiken S.C. 29808

Introduction

With special reference to H diffusion in metals and alloys, diffusion constants, $D_H(c_H)$, determined from flux measurements, J , are a function of the H concentration, c_H , where $D_H(c_H)$ is defined by $J = -D_H(c_H)dc_H/dx$. The concentration-independent diffusion constant, D_H^* , appears in the following equation for the flux, $J = -D_H^*(c_H/RT)(d\mu_H/dx)$ where μ_H is the H chemical potential.

From these two expressions for J , D_H^* can be obtained from D_H , i.e.,

$$D_H = D_H^* f(r) \tag{1}$$

where $f(r)$, the thermodynamic factor, is defined as

$$f(r) = (\partial \mu_H / RT / \partial \ln r)_T = (\partial \ln p^{1/2} / \partial \ln r)_T \tag{2}$$

where p is the H_2 pressure in equilibrium with the H-containing metal phase and $r = (H/metal)$, atom ratio, which is convenient to substitute for c_H in most situations.

Determinations of D_H^* from experimental D_H values have been carried out for Pd-H and Pd-alloy-H systems by Wicke and his students [1]. They employed equation (1) under conditions where r is nearly constant, e.g., they employed a breakthrough method where an increment of H is added to one side of a membrane and the time for the disturbance to reach the other side can be used to calculate D_H . In such an experiment r is essentially constant permitting equation (1) to be employed at the given r value. The thermodynamic factor was determined as a function of r from isotherms for the Pd-H and Pd-alloy-H systems. Küssner [2] determined D_H for the $Pd_{0.77}Ag_{0.23}$ alloy over an H content range from $r=0$ to 0.42 (303 K) and Züchner and Boes [3] carried out similar experiments at 296 K which included the $Pd_{0.60}Ag_{0.40}$ alloy. These alloys do not form hydride phases at these temperatures and therefore they were able to measure D_H over a continuous range of r from which D_H^* values were derived.

Salomons et al [4] measured diffusion constants for Pd from the relaxation time constants for weight changes of Pd samples resulting from sudden pressure changes. D_H values were determined at 473, 513, 563 and 603 K as a function of r up to about 0.08. They found a decrease of $D_H(c_H)$ from $r=0.015$ to ≈ 0.08 and it was explained in terms of the thermodynamic factor times the so-called availability factor, $(1-r)$, which allows for the blocking of interstices by H. They did not measure the dependence of E_D on r .

In this paper the dependences of D_H , D_H° and E_D on r will be discussed from a general point-of-view where these parameters are defined by the Arrhenius equation for the diffusion constant, i.e.,

$$D_H = D_H^\circ \exp(-(E_D/RT)). \quad (3)$$

Results and Discussion

Theoretical Considerations

The experimental dependences of D_H , D_H° and E_D on r will be considered from the viewpoint of a simple model of H in Pd and its alloys valid at small H contents. This is the lattice gas, mean field model given by

$$\mu_H = \mu_H^\circ + RT \ln [(1-r)/r] + \mu_H^E(r) \approx \mu_H^\circ + RT \ln [(1-r)/r] + g_1 r \quad (4)$$

where μ_H is the chemical potential of the dissolved H and μ_H° refers to conditions of $r \rightarrow 0$ without the ideal configurational term and $\mu_H^E(r)$ is the non-ideal or excess chemical potential which, at small r , can be approximated by $g_1 r$ where g_1 is a constant at a given temperature and it is the first order coefficient in a polynomial expansion of $\mu_H^E(r)$ in r [5]. At equilibrium, μ_H in the metal phase must be equal to μ_H in the gas phase, $\frac{1}{2}\mu_{H_2}^\circ + RT \ln p^{1/2}$, and therefore equation (4) can be rewritten as

$$\frac{1}{2}\mu_{H_2}^\circ + RT \ln p^{1/2} = \mu_H = \mu_H^\circ + RT \ln [r/(1-r)] + g_1 r. \quad (5)$$

Since g_1 is a partial free energy, it can be expressed in terms of the partial excess enthalpy and entropy, $g_1 = h_1 - Ts_1$. Values of these parameters for Pd-H have been given by Kuji et al [6] based on experimental data over the range 250-650 K; they are all negative for Pd-H. For example, at 373 K, $g_1 = -46.1$ kJ/mol H, $h_1 = -82.5$ kJ/mol H and $s_1 = -97.5$ J/K mol H [6]. The g_1 values are obtained from slopes of plots of $RT \ln p^{1/2} [(1-r)/r]$ against r and h_1 and s_1 can be derived from the slopes and intercepts of plots of (g_1/T) against $1/T$. The g_1 can be viewed as an H-H interaction free energy which leads to hydride formation below the critical temperature of 565 K [7].

From equations (1), (2) and (5) an expression for the thermodynamic factor, $f(r)$, can be obtained

$$f(r) = 1/(1-r) + g_1 r/RT. \quad (6)$$

From equations (1) and (6), an expression for $\ln D_H$, appropriate for small r , can be derived

$$\ln D_H = \ln D_H^* + \ln f(r) = \ln D_H^* + \ln [1/(1-r) + g_1 r/RT] \approx \ln D_H^* + g_1 r/RT \quad (7)$$

because at small r , $(1/(1-r))$ reduces to 1 and therefore $f(r)$ becomes $g_1 r/RT$.

If the derivative of $\ln D_H$ is taken with respect to $1/T$ in equation (7), we obtain

$$E_D - E_D^* = -h_1 r \quad (8)$$

where E_D^* indicates the concentration-independent E_D . From equation (3) and equations (7) and (8), we obtain

$$\ln (D_H^0/D_H^{0,*}) = -s_1 r / R \quad (9)$$

where $D_H^{0,*}$ is the concentration-independent D_H^0 .

Thus, at small r , the changes of these excess thermodynamic functions with r are given as

$$(d \ln D_H/dr) = g_1/RT; (d E_D/dr)/RT = -h_1/RT; (d \ln D_H^0/dr) = -s_1/R. \quad (10)$$

It is of interest that experimental kinetic parameters, e.g., $(d \ln D_H/dr)$, can be equated to experimental thermodynamic parameters, e.g., g_1/RT .

Experimental Results

The r dependence of D_H and E_D were determined experimentally from H permeation through a Pd membrane. The downstream p_{H_2} was ≈ 0 while the upstream p_{H_2} was maintained at the desired values as determined from the p_{H_2} - r isotherms. For these boundary conditions [8] at relatively small H contents, it can be shown that, $(r_{up}/2) \approx r_{av}$ and the H/Pd values shown in Figure 1 are r_{av} values.

An example of the changes of $\ln D_H$ and E_D with r for diffusion of H at 423 K in the dilute phase of Pd are shown in Figure 1 where it can be seen that the former decreases, while the latter increases, with r as predicted. Their slopes give $g_1 = -37$ kJ/mol H and $h_1 = -91.6$ kJ/mol H which are reasonable for this temperature since a solubility isotherm at 423 K measured here gives $g_1 = -35.9$ kJ/mol H. The value of h_1 from isotherms is not as reliable as the value of g_1 and Kuji et al [5] give $h_1 = -82.5$ kJ/mol H.

It can be concluded that $\ln D_H$ and E_D change with r in predictable ways for the dilute phase of Pd-H. These changes are significant at, e.g., 423 K, E_D increases from about 23.9 ($r=0$) to 25.5 kJ/mol H at $r=0.02$.

Oates and Mohri [9] have used the cluster variation method to describe the Pd-H system concluding that, in addition to the long range non-configurational interaction [10], there is a shorter range non-ideal configurational interaction of the H atoms which is, of course, accompanied by a non-ideal configurational entropy reflected by s_1 . The increase of E_D with r suggests that the formation of the transition state causes some weakening of the H-H attractive interaction. The s_1 term is negative and therefore the non-ideal configuration of the H atoms is partially destroyed by diffusion leading to an increase of the partial excess entropy, $-s_1$.

A more complete account of the dependences of D_H and E_D will be given elsewhere [8]. The strict equating illustrated here of experimental kinetic parameters with thermodynamic parameters, h_1 , g_1 and s_1 , is noteworthy.




Figure 1. H Diffusion parameters for a Pd membrane (75 μm) at 423 K. o, $\ln D_H$
 Δ , E_D .

References

- [1] E. Wicke, H. Brodowsky, in *Hydrogen in Metals, I*, G. Alefeld, J. Volkl, Springer-Verlag, Berlin, 1978.
- [2] A. Küssner, *Z. Naturforschung*, **21a** (1966) 515.
- [3] H. Züchner, N. Boes, *Ber. Bunsenges. Physik. Chem.*, **76** (1972) 783.
- [4] T. Flanagan, W. Oates, *Annu. Rev. Mater. Sci.*, **21** (1991) 269.
- [5] E. Salomons, U. Ljungblad, R. Griessen, *Defect and Diffusion Forum*, **66-69** (1989) 327.
- [6] T. Kuji, W. Oates, B. Bowerman, T. Flanagan, *J. Phys. F:Met. Phys.*, **13** (1983) 1785
- [7] E. Wicke, J. Blaurock, *Ber. Bunsenges. Physik. Chem.*, **85** (1981) 1091.
- [8] T. Flanagan, D. Wang, K. Shanahan, *to be published*.
- [9] T. Mohri, W. Oates, *J. Alloys Compounds*, **330-332** (2002) 14.
- [10] G. Alefeld, *Ber. Bunsenges Physik. Chem.*, **76** (1972) 746.